



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

where V is the velocity of the body in its orbit, k the constant of the system, ψ the angle between the tangent to the new orbit and the prolongation of the corresponding radius vector (r), r_0 the radius vector for the vertex of the old orbit, and p , a , and e are the semi-parameter, semi-major axis, and eccentricity of the new orbit. For the given point we have, as may easily be seen,

$$\begin{aligned} r &= 2m, \\ \psi &= 30^\circ, \\ r^2 \sin^2 \psi &= m^2; \end{aligned}$$

whence, by substitution in (a), we see that the new orbit will be an hyperbola, in which $p = 2m$, $a = -m$, $e = 1\frac{1}{3}$. [Ormond Stone.]



EXERCISES.

26

DETERMINE the maximum right cone B inscribed in a given right cone A , the vertex of B being at the centre of the base of A . [O. Root, Jr.]

27

THE NUMBER of points common to three surfaces of the m th, n th, and p th degrees being in general mnp , find the co-ordinates of all the real and imaginary points of intersection of the three surfaces, $y^2 + x^2 + z^2 = 4x$, $x^2 = yz$, and $y^2 = x^4$. [H. A. Newton.]

28

GIVEN the perpendicular, median, and bisector issuing from one and the same vertex of a plane triangle and terminating in the opposite side, to construct the triangle and determine a formula for its area. [Marcus Baker.]

29

GIVEN a pair of points A, B , if C, D are such that $OD \cdot OC = OA^2 = OB^2$ and $AOC = AOD$, prove that the pair A, B bears similar relation to the pair C, D . Show, also, the existence of a pair E, F which bears the same relation to each of the pairs A, B and C, D . [Wm. Woolsey Johnson.]

30

INVESTIGATE formulæ for the logarithms of 13, 19, and 73 in terms of the logarithms of primes less than 13 and of $(N + 1/N)$ where $N = 132495; 262143; 274625$. [F. H. Loud.]

31

IN THE TRIANGLE ABC draw AD to the point D in BC ; then will

$$AB^2 \cdot DC + AC^2 \cdot BD = BC \cdot AD^2 + BD \cdot DC \cdot BC.$$

[J. R. Spiegel.]

32

A TRIANGLE PQR is inscribed in triangle ABC . Determine the ratios in which P, Q, R divide the sides BC, CA, AB in order that AQR, BRP, CPQ may be respectively $\frac{1}{3}, \frac{1}{5}, \frac{1}{7}$, of ABC . [W. M. Thornton.]

SELECTED.—ALGEBRAIC EQUATIONS.

33

THE CUBIC $ax^3 + 3bx^2 + 3cx + d = 0$ has one or three real roots as its discriminant

$$D = (ad - bc)^2 - 4(b^2 - ac)(c^2 - bd)$$

is positive or negative. If D is zero, two roots are in general equal each to

$$\sqrt{\frac{c^2 - bd}{b^2 - ac}}.$$

34

SOLVE by quadratics the equations below and interpret their roots geometrically:—

$$\begin{aligned} x^4 - x^3 \sqrt{15} + 4x^2 - 1 &= 0, \\ x^4 - x^3 \sqrt{3} - 2x^2 + 2x \sqrt{3} - 1 &= 0. \end{aligned}$$

35

SOLVE by quadratics the quartic

$$x^4 + ax^3 + bx^2 + ad^{\frac{1}{2}}x + d = 0.$$

36

SOLVE the cubic

$$\sqrt[3]{x} + \sqrt[3]{x-7} = \sqrt[3]{3x+3}.$$

by eliminating its commensurable root.

37

IF a, b, c, d, e be the roots of the quintic

$$x^5 + px^3 + qx^2 + rx + s = 0,$$

find the values of

$$a^2(de + eb + bc) + b^2(ea + ac + cd) + c^2(ab + bd + de) \\ + d^2(bc + ce + ea) + e^2(cd + da + ab).$$

38

SHOW that among any three roots a, b, c of a biquadratic, as $x^4 + px^2 + q = 0$, this relations holds:—

$$\frac{1}{a} + \frac{1}{b} + \frac{1}{c} = \frac{1}{a + b + c}.$$

CONICS.

39

ON A CHORD MN of an ellipse as diameter a circle is drawn which cuts the curve in P and Q . Show that the ratio of the intercepts of MN and PQ on the axis major is $e^2 : 2 - e^2$.

40

THE VERTEX of an hyperbola and one asymptote is fixed. Find the locus of the focus.

41

FIND the equations to the conics which cut the ellipse

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$

orthogonally at all their common points.

42

FROM two points on Ox , equidistant from O , tangents are drawn to the conic

$$ax^2 + 2hxy + by^2 = 2x.$$

Find the locus of their points of concurrence.

43

THE POLARS of points on one of two equal circles relative to the other envelop a parabola.

44

FIND the locus of the points from which tangents drawn to $Ax^2 + By^2 = 1$ are parallel to conjugates of $ax^2 + 2hxy + by^2 = 1$.